



G - กึ่งกรุปอันดับที่บรรจุฐานสองด้าน

On Ordered G - Semigroups Containing Two-sided Bases

วิชญาพร จันทะนัน, กชกร นวดไธสง และ สุชาติ เจริญรัมย์*

Wichayaporn Jantanana, Kodchakorn Nuadthaisong and Suchat Jaroenram*

ภาควิชาคณิตศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยราชภัฏบุรีรัมย์

Department of Mathematics, Faculty of Science, Buriram Rajabhat University

Received : 2 September 2020

Revised : 15 December 2020

Accepted : 26 February 2021

บทคัดย่อ

ในบทความวิจัยนี้ศึกษาแนวคิดของ แกมมา กึ่งกรุปอันดับที่บรรจุฐานสองด้าน โดยจะศึกษาค้นคว้าเกี่ยวกับแนวคิดของ แกมมา กึ่งกรุปที่บรรจุฐานสองด้าน ซึ่ง ธวัช ช่างฝัด และ พิสิทธิ์ คำมูล ได้ทำการศึกษาในปี ค.ศ. 2018 นอกจากนี้ เรายังพิสูจน์ว่าทุก ๆ แกมมา กึ่งกรุปอันดับที่บรรจุฐานสองด้านจะมีสมาชิกเท่ากัน

คำสำคัญ : แกมมา กึ่งกรุปอันดับ ; ฐานสองด้าน ; แกมมาไอดีล

Abstract

The aim of this paper is to study the concept of ordered G - semigroups containing two-sided bases that are studied analogously to the concept of G - semigroups containing two-sided bases considered by T. Changpas and P. Kummoon in 2018. Moreover, we prove any ordered G - semigroups containing two-sided bases have the same cardinality.

Keywords : ordered G - semigroup ; two-sided bases ; G - Ideal

*Corresponding author. E-mail : 600112210021@bru.ac.th



Introduction

The notion of two-sided bases of semigroups has been introduced and studied by I. Fabrici. (Fabrici, 1975). Indeed, a non-empty subset A of a semigroup S is said to be a two-sided base of S if A satisfies the two following conditions :

$$(1) S = A \dot{\cup} SA \dot{\cup} AS \dot{\cup} SAS ;$$

$$(2) \text{ If } B \text{ is a subset of } A \text{ such that } S = B \dot{\cup} SB \dot{\cup} BS \dot{\cup} SBS, \text{ then } B = A.$$

The concept of a G -semigroup has been introduced by M. K. Sen. (Sen, 1981). The concept of G -semigroups containing two-sided bases was firstly given by T. Changphas and P. Kummoon. (Thawat & Pisit, 2018). Indeed, a non-empty subset A of S is called a two-sided base of S if it satisfies the two following conditions:

$$(1) S = A \dot{\cup} SGA \dot{\cup} AGS \dot{\cup} SGA GS ;$$

$$(2) \text{ If } B \text{ is a subset of } A \text{ such that } S = B \dot{\cup} SGB \dot{\cup} BGS \dot{\cup} SGB GS, \text{ then } B = A.$$

The main purpose of this paper is to introduce the concept and extend the result to an ordered G -semigroup containing two-sided bases. It will get the form of ordered G -semigroups containing two-sided bases is a non-empty subset A of an ordered G -semigroup S is called a two-sided base of S if it satisfies the two following conditions:

$$(1) S = (A \dot{\cup} SGA \dot{\cup} AGS \dot{\cup} SGA GS) ;$$

$$(2) \text{ If } B \text{ is a subset of } A \text{ such that } S = (B \dot{\cup} SGB \dot{\cup} BGS \dot{\cup} SGB GS), \text{ then } B = A.$$

We now recall some definitions and results used throughout the paper.

Definition 1.1. (Thawat & Pisit, 2018). Let S and G be any two non-empty sets. Then S is called a G -semigroup if there exists a mapping from $S \times G \times S \rightarrow S$, written as $(a, g, b) \mapsto agb$, satisfying the following identity $(aab)bc = aa(bbc)$ for all $a, b, c \in S$ and $a, b \in G$.

Definition 1.2. (Abdul et al., 2017). Let (S, G, ε) be a G -semigroup. For A and B be two non-empty subsets of S , the set product AGB is defined to be the set of all elements agb in S where $a \in A, b \in B$ and $g \in G$. That is

$$AGB := \{agb \mid a \in A, b \in B, g \in G\}.$$

Also we write BGa instead of $BG\{a\}$, and we write aGB instead of $\{a\}GB$, for $a \in S$.

Definition 1.3. (Niovi, 2017). An ordered G -semigroup is a G -semigroup S together with a partial order relation \leq on S such that $a \leq b$ implies $agc \leq bgc$ and $cga \leq cgb$ for all $a, b, c \in S$ and $g \in G$.

(Iampan, 2009). For an element a of an ordered G -semigroup S , define $\langle a \rangle := \{t \in S \mid t \leq a\}$ and for a subset H of S , define $\langle H \rangle = \bigcup_{h \in H} \langle h \rangle$ that is $\langle H \rangle = \{t \in S \mid t \leq h \text{ for some } h \in H\}$. Then the following holds true:



1. $H \cap (H \cup B) = (H \cap B) \cup (H \cap H)$
2. For any subsets A and B of S with $A \subseteq B$, we have $(A \cap C) \subseteq (B \cap C)$;
3. For any subsets A and B of S , we have $(A \cup B) \cap C = (A \cap C) \cup (B \cap C)$;
4. For any subsets A and B of S , we have $(A \cap B) \cap C = (A \cap C) \cap (B \cap C)$.

Definition 1.4. (Niovi, 2017). A non-empty subset A of an ordered G -semigroup (S, G, \leq) is called a G -subsemigroup of S if $AGA \subseteq A$.

Definition 1.5. (Kwon and Lee, 1998). A non-empty subset A of an ordered G -semigroup (S, G, \leq) is called a left (resp. right) G -ideal of S if it satisfies :

- (1) $SGA \subseteq A$ (resp. $AGS \subseteq A$);
- (2) if $a \in A$ and $b \leq a$ for $b \in S$ implies $b \in A$.

Both a left G -ideal and a right G -ideal of an ordered G -semigroup S is called a G -ideal of S .

Definition 1.6. (Kostaq & Edmond, 2006). An G -ideal A of an ordered G -semigroup (S, G, \leq) is called proper if $A \neq S$. A proper G -ideal A of S is called maximal if for each G -ideal T of S such that $A \subseteq T$, we have $T = A$ or $T = S$, i.e., there is no G -ideal T of S such that $A \subseteq T \subseteq S$.

Proposition 1.7. (Kostaq & Edmond, 2006). Let (S, G, \leq) be an ordered G -semigroup and $\{A_i \mid i \in I\}$ a family of G -ideals of S . If $\bigcap \{A_i \mid i \in I\} \neq \emptyset$, then the set $\bigcap \{A_i \mid i \in I\}$ is a G -ideal of S and $\bigcup \{A_i \mid i \in I\}$ is also a G -ideal of S .

It is known (Niovi, 2017) that if denoted by $I(A)$, is the smallest G -ideal of S containing A , and $I(A)$ is of the form $I(A) = (A \cup SGA \cup AGS \cup SGA \cup AGS)$. In particular, for an element $a \in S$, we write $I(\{a\})$, $I(a)$ which is called the principal G -ideal of S generated by a . Thus $I(a) = (a \cup SGA \cup aGS \cup SGA \cup AGS)$. Note that for any $b \in S$, we have $(SGB \cup bGS \cup SGBGS)$ is a G -ideal of S . Finally, if A and B are two G -ideals of S , then the union $A \cup B$ is a G -ideal of S .

Methods

We begin this section with the definitions of two-sided bases of ordered G -semigroups.

Definition 2.1. (Abul *et al.*, 2017). Let (S, G, \leq) be an ordered G -semigroup. A non-empty subset A of S is called a two-sided base of S if it satisfies the two following conditions :

- (1) $S = (A \cup SGA \cup AGS \cup SGA \cup AGS)$;
- (2) If B is a subset of A such that $S = (B \cup SGB \cup BGS \cup SGBGS)$, then $B = A$.

We now provide some examples.



Example 2.2. (Chinnadurai & Arulmozhi, 2018). Let $s = \{a, b, c, d\}$ and $G = \{a, b\}$ where a, b are define on s with the following Cayley tables:

a	a	b	c	d
a	a	a	a	a
b	a	b	c	d
c	a	c	c	c
d	a	c	c	c

b	a	b	c	d
a	a	a	a	a
b	a	b	c	d
c	a	c	c	c
d	a	b	c	d

$$\mathcal{E} := \{(a, a), (a, b), (a, c), (a, d), (b, b), (b, c), (b, d), (c, c), (d, c), (d, d)\}$$

In (Chinnadurai & Arulmozhi, 2018). (s, G, \mathcal{E}) is an ordered G -semigroup. It is easy to see that the two-sided bases of s are $\{b\}$ and $\{d\}$. But $\{b, d\}$ is not a two-sided base.

Example 2.3. (Subrahmanyeswara *et al.*, 2012). Let $s = \{a, b, c, d\}$ and $G = \{a\}$ where a is defined on s with the following Cayley tables:

a	a	b	c	d
a	b	b	d	d
b	b	b	d	d
c	d	d	c	d
d	d	d	d	d

$$\mathcal{E} := \{(a, a), (b, b), (c, c), (d, d), (a, b), (d, b), (d, c)\}$$

In (Subrahmanyeswara *et al.*, 2012). (s, G, \mathcal{E}) is an ordered G -semigroup. It is easy to see that the two-sided bases of s is $\{a, c\}$. But $\{b\}$ and $\{d\}$ are not a two-sided bases.

In Example 2.2. and Example 2.3., it is observed that two-sided bases of s have same cardinality. This leads to prove in Theorem 3.4.

Hereafter, for any ordered G -semigroup (s, G, \mathcal{E}) , we shall use the quasi-ordering which is defined as follows.

Definition 2.4. Let (s, G, \mathcal{E}) be an ordered G -semigroup. We define a quasi-ordering \preceq on s for any $a, b \in s$,

$$a \preceq b \iff I(a) \subseteq I(b).$$

We write $a \prec b$ if $a \preceq b$ but $a \neq b$. It is clear that, for any a, b in s , $a \mathcal{E} b$ implies $a \preceq b$.

Lemma 2.5. Let A be a two-sided base of an ordered G -semigroup (s, G, \mathcal{E}) , and $a, b \in A$.

If $a \in (s G b \mathcal{E} b G s \mathcal{E} s G b G s)$, then $a = b$.



Proof. Assume that $a \in (SGB \dot{\cup} bGS \dot{\cup} SGBGS)$, and suppose that $a \neq b$. Let $B = A \setminus \{a\}$. Since $a \neq b, b \in B$. To show that $I(A) \neq I(B)$, it suffices to show that $A \neq I(B)$. Let $x \in A$. There are two cases to consider. If $x \neq a$, then $x \in B$, and so $x \in I(B)$. If $x = a$, then by assumption we have $x = a \in (SGB \dot{\cup} bGS \dot{\cup} SGBGS) \cap I(B) \cap I(B)$. So we have $I(A) \neq I(B)$. Thus $S = I(A) \neq I(B) \cap S$. This is a contradiction. Hence $a = b$.

Results

In this part, the algebraic structure of an ordered G -semigroup containing two-sided bases will be presented.

Theorem 3.1. A non-empty subset A of an ordered G -semigroup (S, G, \leq) is a two-sided base of S if and only if A satisfies the two following conditions:

- (1) For any $x \in S$ there exists $a \in A$ such that $x \leq a$;
- (2) For any $a, b \in A$, if $a \neq b$, then neither $a \leq b$ nor $b \leq a$.

Proof. Assume first that A is a two-sided base of S . Then $I(A) = S$. Let $x \in S$. Then $x \in I(A) = \bigcup_{a \in A} I(a)$, and so $x \in I(a)$ for some $a \in A$. This implies $I(x) \subseteq I(a)$. Hence $x \leq a$. Hence condition (1) is true. Let $a, b \in A$ such that $a \neq b$. Suppose that $a \leq b$. We set $B = A \setminus \{a\}$. Then $b \in B$. Let $x \in S$. By (1), there exists $c \in A$ such that $x \leq c$. There are two cases to consider. If $c \neq a$, then $c \in B$, thus $I(x) \subseteq I(c) \subseteq I(B)$. Hence $S = I(B)$. This is a contradiction. If $c = a$, then $x \leq a$ hence $x \in I(B)$ since $b \in B$. We have $S = I(B)$. This is a contradiction. The case $b \leq a$ is proved similarly. Hence condition (2) is true.

Conversely, assume that the condition (1) and (2) hold. We will show that A is a two-sided base of S . To show that $S = I(A)$. Let $x \in S$. By (1), then there exists $a \in A$ such that $I(x) \subseteq I(a)$. Then $x \in I(x) \subseteq I(a) \subseteq I(A)$. Thus $S \subseteq I(A)$, and $S = I(A)$. Next it remains to show that A is a minimal subset of S with the property: $S = I(A)$. Suppose that $S = I(B)$ for some $B \subsetneq A$. Since $B \subsetneq A$, there exists $a \in A \setminus B$. So $a \in I(B)$. Since $a \in A \cap I(B) = I(B) \cap I(A)$, it follows that $a \in (SGB \dot{\cup} BGS \dot{\cup} SGBGS)$. Since $a \in (SGB \dot{\cup} BGS \dot{\cup} SGBGS)$, we have $a \leq y$ for some $y \in (SGB \dot{\cup} BGS \dot{\cup} SGBGS)$. There are three cases to consider:

Case 1: $y \in BGS$. Then $y = b_1gs$ for some $b_1 \in B, g \in G$ and $s \in S$. Since $a \leq y$ and $y \in (b_1 \dot{\cup} SGb_1 \dot{\cup} b_1GS \dot{\cup} SGb_1GS)$, we have $a \in (b_1 \dot{\cup} SGb_1 \dot{\cup} b_1GS \dot{\cup} SGb_1GS)$. It follows that $I(a) \subseteq I(b_1)$. Hence, $a \leq b_1$. This is a contradiction.

Case 2: $y \in SGB$. Then $y = sgb_2$ for some $b_2 \in B, g \in G$ and $s \in S$. Since $a \leq y$ and $y \in (b_2 \dot{\cup} SGb_2 \dot{\cup} b_2GS \dot{\cup} SGb_2GS)$, we have $a \in (b_2 \dot{\cup} SGb_2 \dot{\cup} b_2GS \dot{\cup} SGb_2GS)$. It follows that $I(a) \subseteq I(b_2)$. Hence, $a \leq b_2$. This is a contradiction.



Case 3: $y \in SGBGS$. Then $y = s_1g_1b_3g_2s_2$ for some $b_3 \in B, g_1, g_2 \in G$ and $s \in S$. Since $a \notin y$ and $y \in b_3 \in SGb_3 \in b_3GS \in SGb_3GS$, we have $a \in (b_3 \in SGb_3 \in b_3GS \in SGb_3GS)$. It follows that $I(a) \in I(b_3)$. Hence $a \leq b_3$. This is a contradiction.

Therefore A is a two-sided base of S as required, and the proof is completed.

Theorem 3.2. Let A be a two-sided base of an ordered G -semigroup (S, G, \leq) such that $I(a) = I(b)$ for some $a \in A$ and $b \in S$. If $a \neq b$, then S contains at least two two-sided bases.

Proof. Assume that $a \neq b$. Suppose that $b \in A$. Since $a \leq b$ and $a \neq b$, it follows that

$a \in (SGB \in bGS \in SGBGS)$. By Lemma 2.5., we obtain $a = b$. This is a contradiction. Thus $b \in S \setminus A$. Let $B := (A \setminus \{a\}) \cup \{b\}$. Since $b \in B$, we have $b \in A$, and $B \not\subseteq A$. Hence $A \neq B$. We will show that B is a two-sided base of S . To show that B satisfies (1) in Theorem 3.1., let $x \in S$. Since A is a two-sided base of S , there exists $c \in A$ such that $x \leq c$. If $c \neq a$, then $c \in B$. If $c = a$, then $x \leq a$. Since $a \leq b, x \leq a \leq b$, we have $x \leq b$. To show that B satisfies (2) in Theorem 3.1., let $c_1, c_2 \in B$ be such that $c_1 \neq c_2$. We will show that neither $c_1 \leq c_2$ nor $c_2 \leq c_1$. Since $c_1 \in B$ and $c_2 \in B$, we have $c_1 \in A \setminus \{a\}$ or $c_1 = b$ and $c_2 \in A \setminus \{a\}$ or $c_2 = b$.

There are four cases to consider:

Case 1: $c_1 \in A \setminus \{a\}$ and $c_2 \in A \setminus \{a\}$. This implies neither $c_1 \leq c_2$ nor $c_2 \leq c_1$.

Case 2: $c_1 \in A \setminus \{a\}$ and $c_2 = b$. If $c_1 \leq c_2$, then $c_1 \leq b$. Since $b \leq a, c_1 \leq b \leq a$. Thus $c_1 \leq a$, a contradiction. If $c_2 \leq c_1$, then $b \leq c_1$. Since $a \leq b, a \leq b \leq c_1$. So $a \leq c_1$, a contradiction.

Case 3: $c_2 \in A \setminus \{a\}$ and $c_1 = b$. If $c_1 \leq c_2$, then $b \leq c_2$. Since $a \leq b, a \leq b \leq c_2$. Hence $a \leq c_2$, a contradiction. If $c_2 \leq c_1$, then $c_2 \leq b$. Since $b \leq a, c_2 \leq b \leq a$. Thus $c_2 \leq a$, a contradiction.

Case 4: $c_1 = b$ and $c_2 = b$. This is impossible.

Thus B satisfies (1) and (2) in Theorem 3.1. Therefore B is a two-sided base of S .

Corollary 3.3. Let A be a two-sided base of an ordered G -semigroup (S, G, \leq) , and let $a \in A$. If $I(x) = I(a)$ for some $x \in S, x \neq a$, then x belongs to a two-sided base of S , which is different from A .

Theorem 3.4. Let A and B be any two-sided bases of an ordered G -semigroup (S, G, \leq) . Then A and B have the same cardinality.

Proof. Let $a \in A$. Since B is a two-sided base of S , by Theorem 3.1.(1), there exists an element $b \in B$ such that $a \leq b$. Since A is a two-sided base of S , by Theorem 3.1.(1), there exists $a^* \in A$ such that $b \leq a^*$. So $a \leq b \leq a^*$, i.e., $a \leq a^*$. By Theorem 3.1.(2), $a = a^*$. Hence $I(a) = I(b)$. Define a mapping

$$j : A \rightarrow B \text{ by } j(a) = b \text{ for all } a \in A.$$

To show that j is well-defined, let $a_1, a_2 \in A$ be such that $a_1 = a_2, j(a_1) = b_1$ and $j(a_2) = b_2$ for some



$b_1, b_2 \in B$. Then $I(a_1) = I(b_1)$ and $I(a_2) = I(b_2)$. Since $a_1 = a_2$, $I(a_1) = I(a_2)$. Hence $I(a_1) = I(a_2) = I(b_1) = I(b_2)$, i.e., $b_1 \in B$ and $b_2 \in B$. By Theorem 3.1.(2), $b_1 = b_2$. Thus $j(a_1) = j(a_2)$. Therefore, j is well-defined. We will show that j is one- one. Let $a_1, a_2 \in A$ be such that $j(a_1) = j(a_2)$. Since $j(a_1) = j(a_2)$, $j(a_1) = j(a_2) = b$ for some $b \in B$. So $I(a_2) = I(a_1) = I(b)$. Since $I(a_2) = I(a_1)$, $a_1 \in B$ and $a_2 \in B$. This implies $a_1 = a_2$. Therefore, j is one-one. We will show that j is onto. Let $b \in B$. Since A is a two-sided base of S , by Theorem 3.1.(1), there exists an element $a \in A$ such that $b \in A$. Since B is a two-sided base of S , by Theorem 3.1.(1), there exists an element $b^* \in B$ such that $a \in B$. So $b \in A$ and $b^* \in B$, i.e., $b \in B$. This implies $b = b^*$. Hence $I(a) = I(b)$. Thus $j(a) = b$. Therefore, j is onto. This completes the proof.

If a two-sided base A of an ordered G -semigroup (S, G, \mathcal{E}) is a G -ideal of S , then $S = (A \mathcal{E} S G A \mathcal{E} A G S \mathcal{E} S G A G S) \cup A = A$. Hence $S = A$. The converse statement is obvious. Then we conclude that.

Remark 3.5. It is observed that a two-sided base A of an ordered G -semigroup (S, G, \mathcal{E}) is a two-sided G -ideal of S if and only if $A = S$.

In Example 2.2., it is easy to see that $\{d\}$ is a two-sided base of S , but it is not a G -subsemigroup of S . This shows that a two-sided base of an ordered G -semigroup need not to be a G -subsemigroup in (Niovi, 2018). A non-empty subset A of S is called an idempotent if $A = (A G A)$ or $a = a g a$ for all $a \in A$ and $g \in G$. The following theorem gives necessary and sufficient conditions of a two-sided base of S to be a G -subsemigroup S .

Theorem 3.6. A two-sided base A of an ordered G -semigroup (S, G, \mathcal{E}) is a G -subsemigroup if and only if $A = \{a\}$ with $a g a = a$ for all $g \in G$.

Proof. Assume that A is a G -subsemigroup of S . Let $a, b \in A$ and $g \in G$. Since A is a G -subsemigroup of S , $a g b \in A$. Setting $a g b = c$; thus $c \in S G b \in S G b \mathcal{E} b G S \mathcal{E} S G b G B \in (S G b \mathcal{E} b G S \mathcal{E} S G b G B)$. By Lemma 2.5., $c = b$. So $a g b = b$. Similarly, $c \in a G S \in S G a \mathcal{E} a G S \mathcal{E} S G a G S \in (S G a \mathcal{E} a G S \mathcal{E} S G a G S)$. By Lemma 2.5., $c = a$. So $a g b = a$. We have $a = b$. Therefore, $A = \{a\}$ with $a g a = a$ for all $a \in A$ and $g \in G$. The converse statement is clear.

Notation. The union of all two-sided bases of an ordered G -semigroup (S, G, \mathcal{E}) is denoted by R .

Theorem 3.7. Let (S, G, \mathcal{E}) be an ordered G -semigroup. Then $S \setminus R$ is either empty set or a G -ideal of S .

Proof. Assume that $S \setminus R \neq \emptyset$. We will show that $S \setminus R$ is a G -ideal of S . Let $a \in S \setminus R$, $x \in S$ and $g \in G$. To show that $x g a \in S \setminus R$ and $a g x \in S \setminus R$. Suppose that $x g a \notin S \setminus R$. Then $x g a \in R$. Hence $x g a \in A$ for some a two-sided base A of S . We set $b = x g a$ for some $b \in A$. Then $b \in S G a$. By $b \in S G a \in a \mathcal{E} S G a \mathcal{E} a G S \mathcal{E} S G a G S \in (a \mathcal{E} S G a \mathcal{E} a G S \mathcal{E} S G a G S) = I(a)$, it follow that $I(b) \in I(a)$. Next we will show that $I(b) \in I(a)$. Suppose that $I(b) = I(a)$. Since $a \in S \setminus R$ and $b \in A$, $a \neq b$. Since $I(b) = I(a)$ and Corollary 3.3, we conclude



that $a \in R$. This is a contradiction. Thus $I(b) \subseteq I(a)$, i.e., $b \in_p a$. Since A is a two-sided base of s and $a \in s \setminus R$, by Theorem 3.1.(1), there exists $d \in A$ such that $a \in_p d$. Since $b \in_p a \in_p d$, $b \in_p d$. This is a contradiction to the condition (2) of Theorem 3.1., so we have $xga \in s \setminus R$. Similarly, to show that $agx \in s \setminus R$. Suppose that $agx \in R$, then $agx \in A$ for some a two-sided base A of s . Let $agx = c$ for some $c \in A$. Then $c \in aGS$. By $c \in aGS \implies a \in_s Ga \in_s aGS \in_s SGaGS \implies (a \in_s Ga \in_s aGS \in_s SGaGS) = I(a)$, it follow that $I(c) \subseteq I(a)$. Next, we will show that $I(c) \subseteq I(a)$. Suppose that $I(c) = I(a)$. Since $a \in s \setminus R$ and $c \in A$, $a \in_p c$. Since $I(c) = I(a)$ and Corollary 3.3., we conclude that $a \in R$. This is a contradiction. Thus $I(c) \subseteq I(a)$, i.e., $c \in_p a$. Since A is a two-sided base of s and $a \in s \setminus R$, by Theorem 3.1.(1), there exists $e \in A$ such that $a \in_p e$. Since $c \in_p a \in_p e$, $c \in_p e$. This is a contradiction to the condition (2) of Theorem 3.1., so we have $agx \in s \setminus R$. Let $x \in s \setminus R$, $y \in s$ such that $y \notin x$. Next we will show that $y \in s \setminus R$. Suppose that $y \in R$, then $y \in A$ for some a two-sided base A of s . Since A is a two-sided bases of s , by Theorem 3.1.(1) there exists an element $z \in A$ such that $x \in_p z$. Since $y \notin x$, $y \in_p x$ and $x \in_p z$. So we have $y \in_p z$. This is a contradiction. Therefore $y \in R$ then $y \in s \setminus R$. Hence $s \setminus R$ is a G -ideal of s .

Notation. Let M^* be a proper G -ideal of an ordered G -semigroup (s, G, ε) containing every proper G -ideal of s .

Theorem 3.8. Let (s, G, ε) be an ordered G -semigroup and $\mathcal{A} \subseteq R \subseteq s$. The following statements are equivalent:

- (1) $s \setminus R$ is a maximal proper G -ideal of s ;
- (2) For every element $a \in R$, $R \subseteq I(a)$;
- (3) $s \setminus R = M^*$;
- (4) Every two-sided base of s is a one-element base.

Proof. (1) \iff (2). Assume that $s \setminus R$ is a maximal proper G -ideal of s . Let $a \in R$. Suppose that $R \not\subseteq I(a)$. Since $R \not\subseteq I(a)$, there exists $x \in R$ such that $x \notin I(a)$. Hence $x \in s \setminus R$. Since $x \notin I(a)$, $x \in s \setminus R$ and $x \in s$, we have $(s \setminus R) \in I(a) \subseteq s$. Thus $(s \setminus R) \in I(a)$ is a proper G -ideal of s . Hence $s \setminus R \subseteq (s \setminus R) \in I(a)$. This contradicts to the maximality of $s \setminus R$.

Conversely, assume that for every element $a \in R$, $R \subseteq I(a)$. We will show that $s \setminus R$ is a maximal proper G -ideal of s . Since $a \in R$, $a \in s \setminus R$. Hence $s \setminus R \subseteq s$. Since $R \subseteq s$, $s \setminus R \subseteq \mathcal{A}$. By Theorem 3.7., $s \setminus R$ is a proper G -ideal of s . Suppose that M is a proper G -ideal of s such that $s \setminus R \subseteq M \subseteq s$. Since $s \setminus R \subseteq M$, there exists $x \in M$ such that $x \in s \setminus R$, i.e., $x \in R$. Then $x \in M \subseteq R$. So $M \subseteq R \subseteq \mathcal{A}$. Let $c \in M \subseteq R$. Then $c \in M$ and $c \in R$. Since $c \in M$, we have $sGc \subseteq sGM \subseteq M$, $cGs \subseteq MGS \subseteq M$ and $sGcGs \subseteq sGMGS \subseteq M$. Then $I(c) = (c \in_s Gc \in_s cGs \in_s sGcGs) \subseteq M$. Since $c \in R$, by assumption we have $R \subseteq I(c)$. Hence



$S = (S \setminus R) \dot{\cup} R \dot{\cup} (S \setminus R) \dot{\cup} I(c) \dot{\cup} M \dot{\cup} S$. Thus $M = S$. This is a contradiction. Therefore $S \setminus R$ is a maximal proper G -ideal of S .

(3) \hat{U} (4). Assume that $S \setminus R = M^*$. Since $S \setminus R = M^*$, $S \setminus R$ is a maximal proper G -ideal of S . By (1) \hat{U} (2), for every $a \in R, R \in I(a)$. Firstly, we will show that for every $a \in R, S \setminus R \in I(a)$. Suppose that $S \setminus R \not\in I(a)$ for some $a \in R$. Then $I(a) \neq S$. Hence $I(a)$ is a proper G -ideal of S . Thus $I(a) \in M^* = S \setminus R$. Then $I(a) \in S \setminus R$. Since $a \in I(a), a \in S \setminus R$, i.e., $a \in R$. This is a contradiction. Thus $S \setminus R \in I(a)$ for every $a \in R$. Since $S \setminus R \in I(a)$ and $R \in I(a)$ for every $a \in R$, it follows that $S = (S \setminus R) \dot{\cup} R \dot{\cup} I(a) \dot{\cup} I(a) = I(a) \dot{\cup} S$. So $S = I(a)$ for every $a \in R$. Therefore, $\{a\}$ is a two-sided base of S . Let A be a two-sided base of S . We will show that $a = b$ for all $a, b \in A$. Suppose that there exists $a, b \in A$ such that $a \neq b$. Since A is a two-sided base of $S, A \in R$. This is, $a \in R$. So $S = I(a)$. Since $b \in S = I(a)$ and $b \neq a, b \in (S \dot{\cup} a \dot{\cup} S \dot{\cup} S \dot{\cup} a \dot{\cup} S)$. By Lemma 2.5., $a = b$. This is a contradiction. Therefore, every two-sided base of S is an one element base.

Conversely, assume that every two-sided base of S is an one element base. Then $S = I(a)$ for all $a \in R$. We will show that $S \setminus R = M^*$. The statement that $S \setminus R$ is a maximal proper G -ideal of S follows from the proof (1) \hat{U} (2). Let M be a G -ideal of S such that M is not contained in $S \setminus R$. Then $R \subset M \neq \emptyset$. Let $a \in R \subset M$. Hence $a \in R$ and $a \in M$. So $S \dot{\cup} a \in S \dot{\cup} M \in M, a \dot{\cup} S \in M \dot{\cup} S \in M$ and $S \dot{\cup} a \dot{\cup} S \in S \dot{\cup} M \dot{\cup} S$. So we have $I(a) = (a \dot{\cup} S \dot{\cup} a \dot{\cup} S \dot{\cup} a \dot{\cup} S \dot{\cup} S \dot{\cup} a \dot{\cup} S) \in M$. Hence $S = I(a) \in M \in S$. Thus $M = S$. Therefore $S \setminus R = M^*$.

(1) \hat{U} (3). Assume that $S \setminus R$ is a maximal proper G -ideal of S . We will show that $S \setminus R = M^*$. Since $S \setminus R$ is a proper G -ideal of $S, S \setminus R \in M^* \in S$. By assumption, $S \setminus R = M^*$ or $S = M^*$. Since $S \neq M^*$, so we have $S \setminus R = M^*$. The converse statement is obvious.

Discussion

In this research, we investigated the notion of ordered G -semigroup containing two-sided bases. We proved that a non-empty subset A of an ordered G -semigroup (S, G, \leq) is a two-sided base of S if and only if A satisfies the two following conditions (1) for any $x \in S$ there exists $a \in A$ such that $x \leq_l a$; (2) for any $a, b \in A$, if $a \neq b$, then neither $a \leq_l b$ nor $b \leq_l a$. Also, we showed that if A and B be any two-sided bases of an ordered G -semigroup (S, G, \leq) . Then A and B have the same cardinality.

Conclusions

In this research, we have resulted in ordered G -semigroup that is analogously in G -semigroup considered by T. Changpas and P. Kummoon in 2018.



References

- Abul Basar, M.Y. Abbasi and Sabahat Ali Khan. (2017). Some properties of covered \mathcal{G} -ideal in po- \mathcal{G} -semigroups. *International Journal of Pure and Applied Mathematics*, 115(2), 345-352.
- Chinnadurai, V., & Arulmozhi, K. (2018). Characterization of bipolar fuzzy ideal in ordered gamma semigroup. *Journal of the International Mathematical Virtual Institute*, 8, 141-156.
- Fabrics, I. (1975). Two sided bases of semigroups. *Matematicky casopis*, 25(2), 173-178.
- Iampan, A. (2009). Characterizing Ordered Bi-Ideal in Ordered \mathcal{G} -Semigroups. *International Journal of Mathematical Sciences and Informatics*, 4(1), 17-25.
- Kostaq Hila & Edmond Pisha. (2006). Characterizations on ordered \mathcal{G} -semigroups. *International Journal of Pure and Applied Mathematics*, 28(3), 423-439.
- Niovi Kehayopulu. (2011). On regular duo po- \mathcal{G} -semigroup. *Mathematical Institute Slovak Academy of Sciences*, 61(6), 871-884.
- Kwon Y.I. and Lee S.K. (1998). On weakly prime ideals of ordered \mathcal{G} -semigroup. *Comm. Korean Math. Soc.*, 13(2), 251-256.
- Niovi Kehayopulu. (2017). On intra-regular ordered \mathcal{G} -semigroup. *European Journal of Pure and Applied Mathematics*, 10(4), 620-630.
- Sen, M. K. (1981). On \mathcal{G} -semigroup. *Lecture Notes in Pure and Appl. Math.*, 91, 301-308.
- Subrahmanyaswara Rao Seetamraju, V.B., A. Anjaneyulu, A., & Madhusudana Rao, D. (2012). Po- \mathcal{G} -ideals in Po- \mathcal{G} -semigroup. *IOSR Journal of Mathematics*, 1(6), 39-51.
- Thawat Chang phas & Pisit Kummoon. (2018). On \mathcal{G} -Semigroup Containing Two-sided bases. *KKU Sci. J.*, 46(1), 154-161.



Thawat Changphas & Pisit Kummoon. (2018). On left and right bases of a \mathcal{G} -semigroup. *International Journal of Pure and Applied Mathematics*, 118(1), 125-135.